

## Joseph U. Barker, MD www.josephbarkermd.com

## Rehabilitation Protocol: Arthroscopic Meniscectomy/Chondral Debridement

Name:	Date:
Diagnosis:	Date of Surgery:
Phase I (Weeks 0-2)	
Weightbearing: As tolerated with crutches (for balance)	x 24-48 hours – progress to WRAT
• Range of Motion – AAROM → AROM as tolerated	TAZI TO HOUTS Progress to WEITI
Goal: Immediate full range of motion	
Therapeutic Exercises	
<ul> <li>Quad and Hamstring sets</li> </ul>	
<ul><li>Quad and trainstring sets</li><li>Heel slides</li></ul>	
<ul><li>Co-contractions</li></ul>	
<ul> <li>Isometric adduction and abduction exercises</li> </ul>	
<ul> <li>Straight-leg raises</li> </ul>	
<ul><li>o Straight-leg raises</li><li>o Patellar mobilization</li></ul>	
o i atenai modifization	
Phase II (Weeks 2-4)	
<ul> <li>Weightbearing: As tolerated</li> </ul>	
• Range of Motion – maintain full ROM – gentle passive str	retching at end ranges
Therapeutic Exercises	
<ul> <li>Quadriceps and Hamstring strengthening</li> </ul>	
o Lunges	
<ul><li>Wall-sits</li></ul>	
<ul> <li>Balance exercises – Core work</li> </ul>	
_	
Phase III (Weeks 4-6)	
Weightbearing: Full weightbearing	
<ul> <li>Range of Motion – Full/Painless ROM</li> </ul>	
<ul> <li>Therapeutic Exercises</li> </ul>	
<ul><li>Leg press</li></ul>	
<ul> <li>Hamstring curls</li> </ul>	
<ul> <li>Squats</li> </ul>	
<ul> <li>Plyometric exercises</li> </ul>	
<ul> <li>Endurance work</li> </ul>	
<ul> <li>Return to athletic activity as tolerated</li> </ul>	
Comments:	
Frequency: times per week	weeks
Signature:	Date: