

# Joseph U. Barker, MD www.josephbarkermd.com

# Rehabilitation Protocol: Arthroscopic Meniscectomy/Chondral Debridement

Name:	Date:
Diagnosis:	Date of Surgery:

## Phase I (Weeks 0-2)

- Weightbearing: As tolerated with crutches (for balance) x 24-48 hours progress to WBAT
  - Range of Motion AAROM → AROM as tolerated ○ Goal: Immediate full range of motion
- Therapeutic Exercises
  - Quad and Hamstring sets
  - o Heel slides
  - Co-contractions
  - o Isometric adduction and abduction exercises
  - Straight-leg raises
  - Patellar mobilization
- Phase II (Weeks 2-4)
- Weightbearing: As tolerated
- Range of Motion maintain full ROM gentle passive stretching at end ranges
- Therapeutic Exercises
  - Quadriceps and Hamstring strengthening
  - o Lunges
  - Wall-sits
  - Balance exercises Core work

### Phase III (Weeks 4-6)

- Weightbearing: Full weightbearing
- Range of Motion Full/Painless ROM
- Therapeutic Exercises
  - Leg press
  - Hamstring curls
  - Squats
  - Plyometric exercises
  - Endurance work
  - Return to athletic activity as tolerated

### **Comments:**

Frequency:	times per week	Duration:	weeks
requency	unics per week	Durution	Weens

Signature: \_\_\_\_\_

Date: \_\_\_\_\_